

Solution of the Bose-Hubbard model using doorway states

A. N. SALGUEIRO¹, CHI-YONG LIN², A.F.R. DE TOLEDO PIZA¹ and M. WEI-DEMÜLLER³

¹ *Instituto de Física, Universidade de São Paulo, CP 66318 CEP 05389-970, São Paulo, Brazil*

² *Department of Physics, National Dong Hwa University Hua-Lien, Taiwan, R.O.C.*

³ *Physikalisches Institut, Albert-Ludwigs-Universität Freiburg, 79104 Freiburg, Germany*

PACS. 03.75.Lm – Tunneling, Josephson effect, Bose-Einstein condensates in periodic potentials, solitons, vortices, and topological excitations..

PACS. 03.75.Hh – Static properties of condensates; thermodynamical, statistical, and structural properties..

Abstract. – We introduce an efficient method to solve the Bose-Hubbard model. The Schrödinger equation is solved by the successive construction of doorway states. The ground state wavefunction derived by this method contains all relevant many-body correlations introduced by the hamiltonian, but the dimensionality of the Hilbert space is greatly reduced. We apply the doorway method to obtain the chemical potential, the on-site fluctuations and the visibility of the interference pattern arising from atoms in a one-dimensional periodic lattice. Excellent qualitative agreement with exact numerical calculations as well as recent experimental observations is found.

The physics of strongly interacting quantum systems has been subject of intense investigations since the early days of quantum mechanics. With the recent developments in the field of ultracold quantum gases, systems of strongly correlated bosonic and fermionic atoms with tunable interactions have become amenable to experimental studies [1–5]. Complete theoretical understanding of these coupled many-body systems is difficult, and only rare examples exist, for which analytical solutions for the ground state could be explicitly been given [6]. The description generally has to rely upon approximation methods such as the density-group-renormalization [7], a Gutzwiller ansatz [8–10] and quantum Monte Carlo approaches [11]. Of particular importance is the Bose-Hubbard model, which serves as a prototype system exhibiting a quantum phase transition [12]. Triggered by the recent observation of this phase transition from a superfluid to a Mott-insulating phase with bosonic atoms in one- (1D) and three-dimensional (3D) optical lattices [3, 4, 13], the ground state and correlation properties of the Bose-Hubbard model have been extensively investigated by a large number of groups [14–16].

In this Letter, we explore an alternative and particularly efficient theoretical approach to solve the Bose-Hubbard model. The method is based on the construction of doorway projectors as initially introduced by Feshbach in the context of nuclear scattering [17]. A similar approach appears in some other areas of physics under different names. For example, in

solid-state physics it is known as the Lanczos method [18], in quantum optics as the resolvent method [19] and in physical chemistry as the Feshbach-Fano partitioning method [20]. In principle, any state contained in the Hilbert space of the Hamiltonian can serve as a starting state for the doorway method. However, the efficiency of the method can be greatly increased by choosing a start state which is contained in an appropriate subspace determined by the intrinsic symmetries of the hamiltonian. The doorway method thus genuinely leads towards the full many-body wavefunction of the system and it significantly reduces the calculational effort, yet maintaining all correlations of the many-body states.

In this paper, we use the symmetry of the ground-state of the Bose-Hubbard hamiltonian to choose the starting state and to generate the correspondent sequence of doorway states leading into the full wavefunction of the ground state. In order to demonstrate the applicability and the advantages of this method, we determine the energy spectrum of the ground state, which features the phase transition, in both the superfluid and Mott-insulator regime in the homogeneous case. Furthermore, the correlation properties of a bosonic gas in an optical lattice are calculated over the full range of coupling parameters and qualitatively reproduces the measurements of [13,21], even though these experiments are performed in inhomogeneous systems due to the presence of an external trap.

The doorway method consists in splitting the full Hilbert space of the many-body hamiltonian H into different subspaces which are accessed by the action of projectors on H . Let P describe a projector onto a subspace $\mathbb{P}$. For the sake of simplicity, we assume the initial subspace to be one-dimensional with $P = |p\rangle\langle p|$. Note, however, that $\mathbb{P}$ does not necessarily need to be one-dimensional. The first doorway state is given by $|d^{(1)}\rangle \equiv \frac{1}{\mathcal{N}^{(1)}}(1 - P)H|p\rangle$ where $\mathcal{N}^{(1)}$ denotes the normalization, the second doorway by the normalized basis state $|d^{(2)}\rangle \equiv \frac{1}{\mathcal{N}^{(2)}}(1 - P - D_1)H|d^{(1)}\rangle$ and so forth. The procedure terminates when $(1 - P - D_1 - \dots - D_n)H|d^{(n-1)}\rangle = 0$ with $D_i = |d^{(i)}\rangle\langle d^{(i)}|$ ($i = 1, 2, \dots, n$) denoting the projectors onto the doorway spaces. The doorway Lanczos, resolvent or Feshbach-Fano partitioning methods have the same algorithm to generate the so-called doorway states. However, the crucial point is the choice of the most adequate starting state $|p\rangle$ to generate in an efficient way the solution. The space $\mathbb{P} \oplus \mathbb{D}_1 \oplus \dots \oplus \mathbb{D}_n$ then represents an irreducible subspace for the Hamiltonian containing the initial space $\mathbb{P}$. The many-body Schrödinger equation is solved in this subspace

$$(E - H_{PP} - H_{PD_1} \frac{1}{E - H_{D_1 D_1} - M_{1,2}} H_{D_1 P}) P |\psi\rangle = 0, \quad (1)$$

$$M_{i,i+1} \equiv H_{D_i D_{i+1}} (E - H_{D_{i+1} D_{i+1}} - M_{i+1,i+2})^{-1} H_{D_i D_{i+1}}$$

where $H_{PP} = PHP$, $H_{D_i D_j} = D_i H D_j$, $H_{D_i P} = D_i H P$ with $i, j = 1, 2 \dots n$. Recursively, one thus obtains the diagonalization of the Hamiltonian in this subspace. All orders of interactions described by the Hamiltonian are taken into account inside this subspace. Therefore, the expansion is not perturbative in the couplings, but it represents the full solution in a subspace of lower (reduced) dimension.

We apply the doorway method to a system of N bosons in a 1D optical lattice of M sites in the homogeneous case. The system is described by the Bose-Hubbard model $H = -J \sum_{j=1}^M (a_j^\dagger a_{j+1} + h.c.) + \frac{U}{2} \sum_j n_j (n_j - 1)$, which can be seen as a many-mode approximation for an ultracold gas of bosonic atoms occupying the lowest band in a periodic array of potential wells [9,12]. The operators $a_j^\dagger$ and a_j are the creation and annihilation operators, respectively, of a boson in the j -th site, and n_j is the number of atoms at the lattice site j ($j = 1 \dots M$). The strength of the on site repulsion of two atoms on the lattice site i is characterized by U . The parameter J is the hopping matrix element between adjacent sites which characterizes the

strength the tunneling term. A periodic boundary condition is included allowing for tunneling between the first and the last site of the lattice (site $M + 1 \equiv$ site 1).

The Bose-Hubbard hamiltonian with periodic boundary conditions conserves two important quantities: the total number of particles and the total quasi-momentum. As a consequence, its Hilbert space can be partitioned into M decoupled subspaces corresponding to the values of quasi-momentum ($q=0,\dots,M-1$). Furthermore, the subspace with $q=0$ and, for even M , also the subspace with quasi-momentum $q=M/2$, can be further partitioned into symmetric and antisymmetric parts ("odd-even" symmetry) [22]. The ground state of the system lies in the symmetric part of the $q = 0$ subspace. Thus, the initial P subspace is chosen within this manifold. The full set of doorway states constructed from this initial state will finally span the symmetric part of the subspace with zero quasi-momentum, providing the full solution for the ground state. In addition, the ground state of the system has two different limiting cases: the superfluid phase for $J \gg U$ where the particles are delocalized over the whole lattice, and the Mott-insulating phase for $J \ll U$ where the particles are localized with an equal number of atoms per site $n_0 = N/M$.

For the commensurate situation ($N = n_0 M$) we choose $|p\rangle = |n_0\rangle^{\otimes M}$ corresponding to the ground state of the system in the pure Mott case ($U \gg J$). If an extra particle is added to or removed from the system ($N \pm 1 = n_0 M \pm 1$). The appropriate start state is chosen from the symmetric subspace, i.e., $|p\rangle = \frac{1}{\sqrt{M}} \sum_{j=1}^M |(n_0 \pm 1)_j n_0^{\otimes M}\rangle$. Starting from these states, the doorway states are constructed following the method described above. Depending on the parameters of interest, one may also start from the pure superfluid state ($J \gg U$) as the initial state in the construction of the doorways and follow the same procedure as before. Thus, the energy spectrum and eigenvectors are obtained by solving Eq. (1) as a function of the parameters J and U . One can check that the superfluid ground state $|\psi_{SF}\rangle \sim (\sum_{i=1}^M a_i^\dagger)^N |0\rangle$ is asymptotically reached for $J/U \rightarrow \infty$. The doorway method genuinely reduces the numerical effort to the diagonalization of the very small, but relevant part of the total Hilbert space.

A good measure for the convergence of the method is given by the "reliability" defined as $\mathcal{R} = |\langle \psi_0 | \psi_0^{doorway} \rangle|^2 = \sum_{i=0}^n \alpha_i |\langle \psi_0 | d_i \rangle|^2$, where $|\psi_0\rangle$ being the exact ground state of the system obtained from the exact numerical diagonalization of the Bose-Hubbard Hamiltonian, $|\psi_0^{doorway}\rangle$ being the ground state of the system obtained from the doorway method and n being the maximum number of doorways to generate the truncated solution. The meaning of the reliability can be understood in terms of the portion of the full symmetric subspace which has been covered by the solution. The convergency of the method scales linearly with the number of particles for the commensurate and incommensurate cases. Fig.1(a) illustrates this convergency for $M = N = 5$, considering the truncated solution for $n = 2, 3, 4$ and 5 doorways. Excellent convergency is already achieved after the fifth doorway $|d_5\rangle$ is created, showing the subspace spanned by the doorways already include the most relevant portion of the full symmetric subspace. For this particular case, the Hilbert space is of dimension $\binom{N+M-1}{N} = 126$ whereas the symmetric zero quasi-momentum subspace has only dimension 16 and includes the exact ground-state solution. The convergency of the method is very fast, since the recursive construction of the doorway states "automatically" creates the most efficient path into the irreducible subspace depending on the specific values of the parameters U and J . In fact, by applying the doorway method for larger numbers of particles and sites, we find that the number of doorways needed to achieve convergence roughly equals the number of particles.

We derive the ground state energy and the chemical potential ($\mu_N/U = [E_{N+1} - E_N]/U$) as a function of the dimensionless tunneling parameter J/U . Fig. 1(b) shows the chemical potential as a function of J/U comparing the exact solution of the Bose-Hubbard hamiltonian




Fig. 1 – (a) Reliability as a function of the tunneling parameter for $N=M=5$ and the maximum number of doorways $n = 2, 3, 4, 5$. (b) Chemical potential μ_N/U as a function of the dimensionless tunneling parameter J/U for $M = 5$ lattice sites and N particles. The solid lines are the result of the doorway method using seven doorways. The dashed curves represent an exact numerical solution of the Bose-Hubbard model.

(dashed lines) with the result of the doorway method (solid lines) for the case of $M = 5$ sites. The distance between the apparent parallel lines of Fig. 1(b) is almost $1/M$, which vanishes in the thermodynamic limit ($N, M \rightarrow \infty$ and N/M finite), characterizing the phase transition point where the two limiting curves $\mu_{n_0 M}$ and $\mu_{n_0 M+1}$ are the same. For convenience, the doorway method has only been applied to the limiting cases $N = n_0 M$ and $N = n_0 M \pm 1$ where the determination of the corresponding initial states is described above. The curves inside the lobes are only shown for the exact calculation and they represent the evolution of the M -degenerate states of $J = 0$ as a function of the hopping parameter.

As a second application of the doorway method, we determine the on-site number fluctuation of the ground state $\sigma_i = \sqrt{\langle n_i^2 \rangle - \langle n_i \rangle^2}$ as a function of U/J and as a function of J/U . The results are shown in Fig. 2. Depending on the regime, the most efficient construction of the doorways starts from either a pure superfluid state (left graph in Fig. 2) or the Mott state (right graph in Fig. 2). For small U/J , the on-site fluctuations are found to decrease linearly starting from the initial value given by $\sqrt{\frac{N}{M}(1 - \frac{1}{M})}$. At intermediate U/J , the decrease curve changes slope in the region of the phase transition and finally approaches the dependence $2\frac{J}{U}\sqrt{n_0(n_0 + 1)}$, which is identical to the perturbative result for the ground state wavefunction $|\psi\rangle \simeq |n_0\rangle^{\otimes M} + \frac{J}{U} \sum_{\langle i,j \rangle} a_i^\dagger a_j |n_0\rangle^{\otimes M}$ with $\langle i,j \rangle$ denoting the next neighbor sites. The results of the doorway method agree well with an analysis by Roth and Burnett based on the calculation of the superfluid fraction under twisted boundary conditions [14]. Starting from the pure Mott-insulator state, we derive the on-site number fluctuations for $J/U < \sim 1$ (right graph in Fig. 2). In the pure Mott phase, no correlations among the particles exists and the one-site number fluctuations vanish. As J/U increases, correlations are created resulting in a steady increase of the fluctuations. After the phase transition region is crossed, the fluctuations approach the asymptotic value for $U/J = 0$. It is instructive to compare the result with the commonly used Gutzwiller ansatz [9, 10] which, by construction, does not




Fig. 2 – On-site number fluctuations calculated for $M = 6$ sites and $N = 6$ particles using eight doorways to construct the ground state of the system. The initial state in the recursion is either the pure superfluid state (left graph) or the pure Mott-insulator state (right graph). The dashed line gives the result of perturbation theory for $J/U \ll 1$.

provide particle correlations and thus genuinely underestimates on-site fluctuations at small J/U . The short-range correlations of the particles have to be included by additional perturbative terms [10, 16]. Since the groundstate wavefunction constructed by the doorway method contains all relevant many-body correlations, the on-site fluctuations are rendered correctly for small J/U as well as for the region of the phase transition at intermediate values of the tunneling parameter.

To demonstrate the efficiency of the doorway method for the determination of important experimental observables, the interference of atoms released from an optical lattice is modeled and the dependence of the visibility and the width of the first resonance peak of the matter-wave interference pattern on the parameter J/U is derived. Comprehensive studies of these two quantities have recently been performed by Gerbier *et al.* [21, 24] for 3D optical lattice and by Stöferle *et al.* as well as M. Köhl *et al.* for 1D optical lattice [13]. The observed interference pattern is directly related to the Fourier transform of the single-particle density matrix $\rho_{ij} = \langle a_j^\dagger a_i \rangle$. Therefore, measurements of the visibility are particularly sensitive to short-range coherence induced by the tunneling of the particles. Following the derivation in Refs. [14, 25], the intensity is calculated for a one-dimensional lattice as $I = \frac{1}{M} \sum_{i,j=1}^M \exp(ik(r_i - r_j)) \langle a_j^\dagger a_i \rangle$ where k denotes the wave vector of the expanding matter wave after release from the lattice point r_i or r_j , respectively. The result of the doorway method for $N = 8$ particles distributed over $M = 8$ sites, starting from the pure Mott state and taking nine doorways into account, is shown in Fig. 3. Periodic boundary conditions assure that limited size effects are minimized. Fig. 3(a) depicts the resulting intensity distribution for different tunneling parameters. One clearly sees the development of sharp multiple-beam interference maxima with increasing J/U , corresponding to a decrease of the well depth in the experiments. For very small J/U , the remaining weak modulation of the intensity distribution indicates small-scale correlations even deep in the Mott-insulator regime, as have been observed experimentally [13, 21, 24]. The intensity pattern for small J/U is well fitted by the result of perturbation theory for an infinite number of sites given by $I = 1 + 8 \frac{J}{U} \cos(kd)$ (dashed line in Fig. 3) where d is the separation of two adjacent lattice sites.

In Fig. 3(b) the visibility, defined as $V = (I_{\max} - I_{\min}) / (I_{\max} + I_{\min})$, is plotted over a wide range of the parameter U/J . The power of the doorway method becomes apparent by the fact that all features of the visibility are correctly reproduced over the full range of U/J by using a ground state wavefunction created by the superposition of few doorway states including the initial Mott-insulating state. For large U/J (Mott-insulator regime), the visibility approaches the non-vanishing perturbative result $V = 4(n_0 + 1)J/U$ [21, 24, 26] reflecting the persistence of short-range correlations. In the ballpark of the phase transition the visibility changes its analytical dependence on J/U and finally reaches a value close to unity for small U/J . This graph qualitatively describes the experimental findings of Gerbier *et al.* [21], even though the present model neglects the external trap and it is not 1D. A detailed investigation of the visibility in a 3D- inhomogeneous case can be found at ref. [27]. In addition, we can directly calculate the width of the first interference peak for different number of sites and compare it with the experimental measurements of refs. [13] for 1D optical lattice. In Fig. 3(c), the width of the first interference peak is calculated using the doorway method for different number of sites ($M = 3, 4, 5, 6, 7, 8$). We observe that the peak width starts to increase at a critical value close to $U/J \sim 4$ which characterizes the phase transition region. Our calculation describes qualitatively the results of [13] close to the phase transition point. At larger $U/J > 10$ the width of the interference peak in our model is limited by the finite number of sites.

In conclusion, we introduce an alternative and powerful theoretical method to solve the Bose-Hubbard model for interacting bosonic particle in a periodic lattice. The method is based on the successive construction of doorway states which genuinely takes profit of symmetries included in the hamiltonian. It naturally terminates when the full irreducible subspace containing the starting state is spanned by the doorway states. The intermediate doorway states depend on the interaction and tunneling parameters of Bose-Hubbard hamiltonian providing the closest approximation to the ground state. Therefore, the convergence of the method, as can be quantified by the “reliability” introduced in this paper, scales as the number of particles. All relevant correlations of the many-body system are included in the ground state wavefunction derived in this way, as was shown by calculating the energy spectrum, the on-site fluctuations and the expectation value for single-particle correlations. We have limited our discussion on a one-dimensional model to keep the calculational effort on a low level. Conceptually, there are no limitations for extending the method to 2D, 3D and to the inhomogeneous case. For a more detailed comparison with experimental findings, in particular concerning the effects of an additional trapping potential [27], we will also add a harmonic term in the Hamiltonian leading to the appearance of different phases within the sample. Other important observables currently under experimental investigation, such as two-particle correlations of Hanbury-Brown and Twiss type [28], can also be approached by the doorway method. Since the doorway method allows direct access to the many-body wavefunction of the system, it can be applied to other important systems involving many-body correlations such as fermionic gases coupled by Feshbach resonances or the fermionic variant of the Bose-Hubbard model. In addition, dynamical properties of the Bose-Hubbard model, such as many-body tunneling rates, may be investigated.

* * *

We thank W. Zwerger, A. Buchleitner, A. Malvezzi, T. Stöferle and T. Esslinger for enlightening discussions. In particular we thank T. Stöferle and T. Esslinger to provide us with their experimental data. A.N.S. is grateful to M. da Mata and is financially supported by FAPESP. M.W. acknowledges support by the DAAD in the framework of the PROBRAL programme. M.W. and A.N.S. thank L.G. Marcassa and V.S. Bagnato (USP São Carlos)




Fig. 3 – (a) Interference pattern of a matter waves released from a one-dimensional lattice for different values of the tunneling parameter J/U for eight particles distributed over eight lattice sites. The intensity is plotted as a function of the accumulated difference in phase after the expansion. (b) Visibility of the interference pattern as a function of the inverse tunneling parameter U/J . (c) The width of the first interference peak as a function of U/J for different number of sites $M = 4, 5, 6, 7, 8$. The dashed lines in (a) and (b) show the result of perturbation theory for an infinite number of sites but finite average population per site $n_0 = 1$.

for their hospitality. The work of C-Y.L. was partially supported by the National Science Council, ROC under the Grant NSC-94-2112-M-259-008 and by FAPESP. C-Y.L. thanks the hospitality of the members of Departamento de Física Matemática of USP, where this work was performed.

REFERENCES

- [1] M. W. ZWIERLEIN *et al.*, *Nature*, **435** (2005) 1047.
- [2] C. CHIN *et al.*, *Science*, **305** (2004) 1128.
- [3] B. P. ANDERSON E M. A. KASEVICH, *Science*, **282** (1998) 1686.
- [4] M. GREINER *et al.*, *Nature*, **515** (2002) 39.
- [5] MICHAEL KÖHL *et al.*, *Phys. Rev. Lett.*, **94** (2005) 080403.
- [6] E. H. LIEB AND W. LINIGER, *Phys. Rev.*, **130** (1963) 1605.

- [7] S. R. WHITE, *Phys. Rev. Lett.*, **69** (1992) 2863; T. D. KÜHNER AND H. MONIEN, *Phys. Rev. B*, **58** (1998) R14741; S. RAPSCH, U. SCHOLLWÖCK AND W. ZWERGER, *Europhys. Lett.*, **46** (1999) 559 and references therein.
- [8] D.S. ROKSHAR AND B.G. KOTLIAR, *Phys. Rev. B*, **44** (1991) 10328.
- [9] D. JAKSCH *et al.*, *Phys. Rev. Lett.*, **81** (1998) 3108.
- [10] C. SCHROLL, FLORIAN MARQUARDT AND C. BRUDER, *Phys. Rev. A*, **70** (2004) 053609, and references therein.
- [11] G. G. BATROUNI *et al.*, *Phys. Rev. Lett.*, **89** (2002) 117203 and references therein;
- [12] M. FISHER *et al.*, *Phys. Rev. B*, **40** (1989) 546.
- [13] THILO STÖFERLE *et al.*, *Phys. Rev. Lett.*, **92** (2004) 130403; M. KÖHL *et al.*, *Appl. Phys. B*, **79** (2004) 1009.
- [14] R. ROTH AND K. BURNETT, *Phys. Rev. A*, **67** (2003) 031602(R).
- [15] R. ROTH AND K. BURNETT, *Phys. Rev. A*, **68** (2003) 023604.
- [16] W. KRAUTH, M. CAKAREL, AND J.-P. BOUCHARD, *Phys. Rev. B*, **45** (1992) 3137; J.K. FREERICKS AND H. MONIEN, *Phys. Rev. B*, **54** (1996) 16172; D. VAN OOSTEN, P. VAN DER STRATEN AND H. T. C. STOOF, *Phys. Rev. A*, **63** (2001) 053601; W. ZWERGER, *Journal Of Optics B: Quantum Semiclass. Opt.*, **5** (2003) S9-S16; H. P. BÜCHLER *et al.*, *Phys. Rev. Lett.*, **90** (2003) 130401; A. F. R. DE TOLEDO PIZA, *Brazilian Journal of Physics*, **34** (2004) 1102.
- [17] H. FESHBACH, A.K. KERMAN AND R.H. LEMMER, *Annals of Physics*, **41** (1967) 230.
- [18] A good review about Lanczos algorithm can be found at A. MALVEZZI, *Braz. J. Phys*, **33** (2003) 55 and references therein; G. GIUSEPPE GROSSO AND G. PASTORI PARRAVIVINI, *Solid State Physics*, edited by ED. ACADEMIC PRESS 2002 and references therein.
- [19] CLAUDE COHEN-TANNOUDJI, JACQUES DUPONT-ROC, GILBERT GRYNBERG, *Photons and Atoms: Introduction to Quantum Electrodynamics*, edited by ED. WILEY-INTERSCIENCE 1989.
- [20] V. BREMS *et al.*, *J. Chem. Phys.*, **117** (2002) 10635; V. BREMS *et al.*, *J. Chem. Phys.*, **116** (2002) 8318; V. BREMS *et al.*, *J. Chem. Phys.*, **104** (1996) 2222; V. BREMS *et al.*, *J. Chem. Phys.*, **103** (1995) 4524.
- [21] F. GERBIER *et al.*, *Phys. Rev. Lett.*, **95** (2005) 050404.
- [22] A. R. KOLOVSKY AND A. BUCHLEITNER, *Europhys. Lett.*, **68** (2004) 632.
- [23] J. K. FREERICKS AND H. MONIEN, *Phys. Rev. B*, **53** (1996) 2691.
- [24] F. GERBIER *et al.*, *Phys. Rev. A*, **72** (2005) 053606.
- [25] V. A. KASHURNIKOV, N. V. PROKOF'EV, AND B. V. SVISTUNOV, *Phys. Rev. A*, **66** (2002) 031601(R).
- [26] Y. YU, *Phys. Rev. A*, **72** (2005) 053629
- [27] C. KOLLATH *et al.*, *Phys. Rev. A*, **69** (2004) R031601; P. SENGUPTA *et al.*, *Phys. Rev. Lett.*, **95** (2005) 220402.
- [28] SIMON FÖLLING *et al.*, *Nature*, **434** (2005) 481.